



ELSEVIER

JAMDA

journal homepage: www.jamda.com

Special Article

Recommendations on Physical Activity and Exercise for Older Adults Living in Long-Term Care Facilities: A Taskforce Report

Philippe de Souto Barreto PhD^{a,b,*}, John E. Morley MB, BCh^c,
 Wojtek Chodsko-Zajko PhD^d, Kaisu H. Pitkala MD, PhD^e,
 Elizabeth Weening-Dijksterhuis PT, PhD^f, Leocadio Rodriguez-Mañas MD, PhD^g,
 Mario Barbagallo MD, PhD^{h,i}, Erik Rosendahl PT, PhD^j, Alan Sinclair MD^k,
 Francesco Landi MD^l, Mikel Izquierdo PhD^m, Bruno Vellas MD, PhD^{a,b},
 Yves Rolland MD, PhD^{a,b} under the auspices of The International Association of
 Gerontology and Geriatrics – Global Aging Research Network (IAGG-GARN) and the
 IAGG European Region Clinical Section

^a Gerontopole of Toulouse, University Hospital of Toulouse (CHU-Toulouse), Toulouse, France

^b UMR INSERM 1027, University of Toulouse III, Toulouse, France

^c Divisions of Geriatric Medicine and Endocrinology, Saint Louis University School of Medicine, St Louis, MO

^d Graduate College, University of Illinois at Urbana-Champaign, Champaign, IL

^e Unit of Primary Health Care, Department of General Practice and Helsinki University Hospital, University of Helsinki, Helsinki, Finland

^f Lectoraat Healthy Aging, Allied Health Care and Nursing, School of Health Care Studies, Hanze University, Groningen, the Netherlands

^g Service of Geriatrics, Getafe University Hospital, Madrid, Spain

^h International Association of Gerontology and Geriatrics for the European Region, Chair of the Clinical Section, Palermo, Italy

ⁱ University of Palermo, Palermo, Italy

^j Department of Community Medicine and Rehabilitation, Physiotherapy, Umeå University, Umeå, Sweden

^k University of Aston & Diabetes Frail, Birmingham, United Kingdom

^l Department of Geriatrics, Neurosciences and Orthopedics, Catholic University of the Sacred Heart, Rome, Italy

^m Department of Health Sciences, Public University of Navarre, Navarra, Pamplona, Spain

A B S T R A C T

Keywords:

Physical activity
 exercise
 elderly
 long-term care
 nursing home
 functional ability

A taskforce, under the auspices of The International Association of Gerontology and Geriatrics—Global Aging Research Network (IAGG-GARN) and the IAGG European Region Clinical Section, composed of experts from the fields of exercise science and geriatrics, met in Toulouse, in December 2015, with the aim of establishing recommended recommendations of physical activity and exercise for older adults living in long-term care facilities (LTCFs). Due to the high heterogeneity in terms of functional ability and cognitive function that characterizes older adults living in LTCFs, taskforce members established 2 sets of recommendations: recommendations for reducing sedentary behaviors for all LTCF residents and recommendations for defining specific, evidence-based guidelines for exercise training for subgroups of LTCF residents. To promote a successful implementation of recommendations, taskforce experts highlighted the importance of promoting residents' motivation and pleasure, the key factors that can be increased when taking into account residents' desires, preferences, beliefs, and attitudes toward physical activity and exercise. The importance of organizational factors related to LTCFs and health care systems were recognized by the experts. In conclusion, this taskforce report proposes standards for the elaboration of strategies to increase physical activity as well as to prescribe exercise programs for older adults living in LTCFs. This report should be used as a guide for professionals working in LTCF settings.

© 2016 AMDA – The Society for Post-Acute and Long-Term Care Medicine.

The authors declare no conflicts of interest.

* Address correspondence to Philippe de Souto Barreto, PhD, Gerontopole of Toulouse, Institute of Ageing, 37, Allées Jules Guesde, 31000 Toulouse, France.

E-mail address: philipebarreto81@yahoo.com.br (P. de Souto Barreto).

<http://dx.doi.org/10.1016/j.jamda.2016.01.021>

1525-8610/© 2016 AMDA – The Society for Post-Acute and Long-Term Care Medicine.

Older adults living in long-term care facilities (LTCFs) are a complex population, being characterized by high prevalence of dependency in activities of daily living (ADLs), multimorbidity, and

polymedication.¹ Providing the best care for this population represents an immense challenge, particularly in the context of demographic projections for the coming decades in terms of the population aging. In its recent report,² the United Nations estimated that the number of people 60 years or older is expected to more than double between 2013 and 2050, with people 80 years or older constituting the age group with the fastest rate of growth. The number of people living in LTCFs^{3,4} is also expected to rise, leading to an important increase in health care costs.^{3–6}

One of the key challenges for the care of the institutionalized elderly is to maintain residents' functional ability, which is made up of subjects' intrinsic capacity and environmental characteristics,⁷ and the ability to cope with their functional limitations for as long as possible. Overall physical activity has been shown to protect against both the incidence of ADL disability and progression of the disability severity in the general population.⁸ Experts in LTCF research and clinical care, with the support of the International Association of Gerontology and Geriatrics (IAGG) and the World Health Organization, have already recognized the importance of exercise for the quality of care in the LTCF setting.⁹ Scientific evidence from recent meta-analyses have shown that exercise training (ie, a subset of physical activity that is planned, structured, repetitive, and purposeful, being generally used to improve/maintain physical and functional capacities), has been found to have positive effects on the ability to perform ADLs in LTCF residents.^{10,11} Exercise training presents undoubtable advantages because it can positively impact several clinical outcomes that are often present in LTCF residents (eg, falls, cardiovascular diseases, mood disorders), with a low risk of adverse health events. Although exercise undeniably provides health benefits for older people,¹² current physical activity guidelines for older adults (see Table 1) were established from a public health perspective (mainly focusing on the primary prevention of noncommunicable diseases).^{13,14} Due to its specificities (functional limitations, multimorbidity), the exercise-related objectives for LTCF residents understandably focus more on the maintenance of functional ability and improvements in quality of life than the primary prevention of noncommunicable diseases. Therefore, the current physical activity guidelines for older adults are certainly more appropriate for healthy community-dwelling older adults than to very old and vulnerable institutionalized elderly.

The present article reports on the results of a taskforce held in Toulouse, France, on December 1, 2015, before the Nursing Home Research International Working Group Conference (Toulouse, December 2 and 3, 2015), under the auspices of the IAGG–Global Aging Research Network (IAGG–GARN) and the IAGG European Region Clinical Section. The main objectives of this taskforce, which involved experts from the broad fields of geriatrics, particularly nursing home care, and exercise and sports sciences, particularly physiotherapy and exercise for geriatric populations, were to define strategies to increase physical activity and to establish exercise guidelines for people living in LTCFs. Scientific evidence and feasibility issues for the implementation of physical activity strategies and exercise interventions in a long-term basis in LTCFs were the main pillars for the elaboration of the guidelines reported hereafter.

Health Care Issues in LTCFs and Potential Exercise Benefits

Besides ADL dependency, LTCF residents face other important medical challenges. Dementia care, behavioral and psychological symptoms of dementia (BPSD), falls, malnutrition, pain, the use of potentially harmful drugs (eg, antipsychotics), and mood (particularly depression), being severely sedentary (bed and chair rest), and quality of life are often recognized by LTCF staff and experts as crucial issues for the care of residents.^{15,16} Exercise training has the potential to improve many of these issues.

ADL Performance

The most robust information about the positive effects of exercise for people in LTCFs comes from the review and meta-analysis of randomized controlled trials (RCTs) from the Cochrane Group.¹⁰ In this review, Crocker et al¹⁰ found that those who exercised had better ability to perform ADLs (as measured by the Barthel Index or by the Rivermead mobility index) than controls; when pooling the results of all RCTs regardless of the tools used to measure ADLs, Crocker et al¹¹ found a positive effect of exercise (standardized mean difference [SMD] 0.24, 95% confidence interval [CI] 0.11–0.38; $P = .0005$; 13 studies, 2363 participants), roughly corresponding to an increase of 1.3 points on the Barthel Index 20-point scale; such improvement means, for example, that a person who was dependent on bathing became independent.

Table 1
Current Physical Activity Guidelines for Older Adults^{12,13}

Minimum Recommendations of Physical Activity for Adults 65 Years and Older	
<i>Recommendation from the American College of Sports Medicine and the American Heart Association (2007)*</i>	<i>Recommendation from the World Health Organization (2010)[†]</i>
Aerobic/endorurance (bouts of >10 min). Moderate-intensity aerobic PA, 30 minutes, 5 d/wk or vigorous-intensity aerobic PA, 20 min, 3 d/wk. These moderate or vigorous PAs are in addition to the light-intensity activities performed in daily life (eg, self-care) or moderate-intensity PAs lasting 10 min or less	Aerobic/endorurance (bouts of ≥10 min). Moderate-intensity aerobic PA, ≥ 150 min/wk or vigorous-intensity aerobic PA, ≥ 75 min/wk
+	+
Strength. 8–10 exercises, 10–15 repetitions, for strengthening the major muscles of the body, ≥ twice/wk, moderate to high intensity	Strength. ≥ twice/wk, muscle-strengthening involving major muscle groups
+	+
Flexibility. ≥ twice/wk, 10 min	Balance/coordination. ≥ 3 d/wk, for older adults with poor mobility in order to enhance balance and prevent falls.
+	
Balance/coordination. "To reduce risk of injury from falls, community-dwelling older adults with substantial risk of falls should perform exercises that maintain or improve balance." ¹²	

PA, physical activity.

*Adapted from: Nelson ME, Rejeski WJ, Blair SN, et al. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39:1435–1445.

[†]Adapted from: World Health Organization. *Global Recommendations on Physical Activity for Health*. Geneva: WHO Press, 2010.

Dementia Care and Cognitive Function

Recent systematic reviews suggest that exercise is of benefit for the mobility and physical function of people with dementia.^{17,18} Forbes et al,¹⁷ in a meta-analysis of RCTs on exercise for people with dementia, found that exercisers had better ADL than controls (SMD 0.68, 95% CI 0.08 to 1.27, $P = .03$; 6 trials, 289 participants), even though the authors judged this result to be of low quality (due to inconsistency and imprecision). Although the study by Forbes et al¹⁷ was not restricted to LTCF residents, the large majority of the included trials were performed among institutionalized older adults (only 2 of 17 RCTs were developed among community-dwellers). Mixed results have been found for a potential positive impact of exercise on cognitive function in LTCF residents,^{10,17,18} with no clear and robust conclusions established currently.

BPSD and Depression

Very recent findings from a meta-analysis of RCTs on exercise for treating BPSD in people with dementia did not find a statistically significant effect of exercise on the global levels of BPSD.¹⁹ However, the authors found a positive effect of exercise on depressive symptoms, including when restricting analysis to people living in LTCFs (SMD -0.323 [-0.628 to -0.018], $P = .038$; 6 trials, 427 participants); exploratory analysis showed that exercise significantly reduced aberrant motor behavior and provided promising results (not statistically significant) for reducing apathy, agitation, and eating disorders. Although that meta-analysis found a significant effect of exercise on depressive symptoms for people with dementia in LTCFs, the literature on this topic is still mixed. Indeed, although some RCTs in LTCFs found that exercise has reduced depressive symptoms,^{20–22} others found no effect,^{23–28} and one study showed that exercise increased depressive levels in a subpopulation of people with impaired cognition.²⁹ One of the largest RCTs of exercise designed to reduce depression in residents of care homes³⁰ found that the intervention (twice per week, 45 minutes, physiotherapist-led group exercise sessions plus a whole home component designed to encourage more physical activity in daily life) was not effective in reducing both depressive symptoms and clinical depression in a 1-year interval; it is important to highlight that participants' compliance in this study was low (approximately 50% of exercise sessions, which means that participants really performed exercises on average only once a week).

Falls

Evidence for falls prevention in LTCF through exercise is still mixed. The meta-analysis of RCTs by Cameron and colleagues³¹ found no significant effect of exercise (8 RCTs, 1844 participants) on both the rate of falls and the number of fallers. Although the authors suggested that exercise appeared to reduce falls in less frail individuals and to increase falls in frailer people, these results were not significant and were obtained after performing subgroup analysis in a small number of studies (4 RCTs for each of "less frail" and "frailer"). Although it is not possible to crudely translate the findings obtained in other populations/settings to the LTCF population, the results of a meta-analysis of RCTs on falls in community-dwelling older adults,³² which included many more RCTs (59 exercise RCTs, 13,264 randomized participants) than the study by Cameron et al,³¹ found that multicomponent exercise (a combination of 2 exercise types or more among the following: balance/functional training, strength, endurance, flexibility, 3D [exercises such as Tai Chi], general physical activity, or others) reduced both the rate of falls and the risk of falling. Moreover, this study³² and another meta-analysis of RCTs on fall-induced injuries in older community-dwellers,³³ showed that exercise decreased the risk of fractures and injurious falls (including severely injurious falls). Similarly, a

review³⁴ of exercise RCTs in older adults with physical frailty, a very common condition in LTCFs,^{35,36} showed that 7 of 10 studies investigating the effects of exercise on incident falls found low rates of falls among exercisers compared with controls.

Quality of Life

Evidence for quality of life is very sparse,³⁷ with mixed results across RCTs in the institutionalized elderly. Whereas Lee et al³⁸ found improvements in health-related quality of life (both physical and mental components) in a Tai Chi group and Mihalko et al³⁹ showed improvements in life satisfaction in a strength exercise group compared with controls, Kerse et al,²⁹ McMurdo and Rennie,²² and Conradsson et al⁴⁰ obtained no differences between exercisers and controls after the exercise intervention, although this latter found a positive effect of exercise on subjective well-being in the subgroup of people with dementia at 3 months; Chin A Paw et al²³ found slight reductions in quality of life among exercisers compared with controls. Although on the basis of the scientific evidence currently available it is impossible to draw any solid conclusions, the positive impact of exercise on key health aspects for older populations, such as improvements on functional ability,¹⁰ physical function (including the ability to walk),^{10,41} and depression,^{19,42} suggests that exercise should improve quality of life in older people in LTCFs.

Bed and Chair Rest and Sarcopenia

Bed and chair rest are important issues in LTCFs, with residents spending approximately 75% of their waking time in sedentary activities.^{43,44} The use of both psychotropic drugs and physical restraints contribute to the high levels of physical inactivity in LTCF residents. Bed and chair rest may lead to muscle disuse, which, in turn, increases the risk of sarcopenia.^{45,46} Exercise may contribute to decreasing the risk of sarcopenia, which is very prevalent in the institutionalized elderly.^{47–50} Indeed, because sarcopenia is defined by low muscle mass and low muscle strength/functional impairment,^{47–51} exercise could decrease sarcopenia in LTCFs by improving muscle strength and overall physical function. Exercise is a well-known powerful intervention for improving strength and physical function in older adults,⁵² including those living in LTCFs.¹⁰ For instance, Crocker et al¹⁰ showed positive effects of exercise on gait speed in LTCF residents; moreover, most RCTs included in the meta-analysis by Crocker et al¹⁰ found positive effects of exercise on strength, particularly in the lower body. Although it is unlikely that exercise alone has positive effects on anorexia, weight loss, or the overall nutritional status of residents, all of them being common clinical conditions in LTCFs that lead to or increase the severity of sarcopenia, some evidence supports the conclusion that exercise plus nutritional supplementation improves the nutritional status of frail older adults.⁵³

Drugs

Data on the impact of exercise in decreasing the use of harmful drugs, such as antipsychotics, is almost nonexistent. In the meta-analysis by de Souto Barreto et al,¹⁹ only 4 RCTs (3 of them in LTCFs) of 20 reported data on changes in antipsychotic use, with no changes in 3 of them, and mixed results in a fourth RCT⁴⁰; none of the 4 RCTs reported data on the doses of antipsychotics. In a small RCT, Landi et al⁵⁴ showed that exercise training reduced the use of antipsychotics and hypnotics in cognitively impaired people.

Besides its importance in the treatment/prevention of some of the clinical issues previously mentioned, exercise is a recommended intervention for the treatment of several diseases that are prevalent in LTCF residents,^{55,56} such as hypertension,⁵⁷ heart diseases,^{58,59} osteoarthritis,⁶⁰ osteoporosis,^{61,62} diabetes,⁶³ and stroke.⁶⁴

In sum, the current evidence clearly shows that exercise delays age-related declines in the ability to perform ADLs and improves physical function in older people living in LTCFs. Although exercise has the potential to improve other health outcomes in this population, such as falls, quality of life, BPSD, and depression, the results of RCTs have been mixed and the findings are, therefore, still inconsistent.

Physical Activity and Exercise Recommendations for Subgroups of the LTCF Population

Exercise may improve the health and daily care of institutionalized older populations; however, older adults living in LTCFs constitute a heterogeneous population. In terms of functional ability, residents may transition from independent living, to assisted living, and finally into skilled nursing care. Similarly for cognitive functioning, residents often transition from intact cognitive function, to mild cognitive impairment, and to dementia with various degrees of severity (from mild to severe dementia). This heterogeneity suggests that a single exercise prescription is unlikely to fit the desires and needs of all LTCF residents.

Taskforce members decided, therefore, to organize their recommendations into 2 different levels: a first level of recommendations aiming at reducing sedentary behaviors for all LTCF residents, and a second-level set of recommendations that aims to establish relatively specific, evidence-based guidelines for exercise training for well-defined subgroups of LTCF residents (Box 1).

Improving Overall Daily Life Levels of Activity Among all LTCF Residents: First-Level of Recommendations

Although, to the best of our knowledge, no data are available on the negative role of sedentary behaviors for the health of LTCF residents, limited evidence from community-dwelling individuals suggests that sedentary time (ie, waking time spent in activities that did not increase energy expenditure over 1.5 metabolic equivalents,⁶⁵ such as

watching TV) is associated with adverse health outcomes, including the incidence of both type 2 diabetes and cardiovascular disease, and mortality.^{66–68} Interestingly, the associations between sedentary time and adverse health events are independent of subjects' physical activity levels. In a meta-analysis, Chau et al⁶⁸ found a 34% higher mortality risk for adults sitting 10 hours per day. Because LTCF residents spent approximately 75% of their waking time^{43,44} or more than 12 hours per day⁶⁹ in sedentary activities, they constitute a target population that might benefit from interventions designed to reduce sedentary time.⁷⁰

The increase of daily life activity levels in LTCFs is related to the reduction of sedentary time. In the LTCF setting, it is probably feasible and appropriate to elaborate strategies aiming at reducing sitting time through an increase in light physical activities.⁷¹ For example, Slaughter and Estabrooks,⁷² in a preliminary, quasi-experimental study showed that asking LTCF residents to stand up and sit down as many times as possible once in the morning and once in the evening improved residents' functional fitness as measured by the 30-second chair rise test.

Motivation and pleasure are the key aspects to take into account when attempting to increase overall activity levels in such a population, in whom ADL disability and cognitive impairment are common. To increase residents' motivation, it is important to build awareness of the importance of replacing sedentary time with physically demanding activities, even if those activities are of light intensity (eg, walking slowly, light gardening), and LTCF staff should promote residents' physical engagement during social and daily life activities. Building awareness should target both the older residents themselves as well as LTCF staff, other health care professionals (including the primary care physician), residents' family, and policy makers. To increase motivation and impact overall activity levels, the needs of both residents and LTCF staff must be assessed. This assessment of needs will permit LTCF staff to understand potential motivators and barriers to increasing resident activity levels, as well as insight into some of the motivators and barriers related to LTCF staff implementing the appropriate intervention strategies.^{73,74}

Box 1. Summary of Key Recommendations of Physical Activity and Exercise for Older Adults in Long-Term Care Facilities

Increasing Overall Physical Activity Levels in Daily Life	Exercise Training for Residents Dependent in Basic Activities of Daily Living, but Capable of Ambulating/Rising From a Chair*
<ol style="list-style-type: none"> Motivation and pleasure are the key aspects to take into account so as to increase overall activity levels among older adults living in long-term care facilities (LTCFs). LTCF staff should adopt strategies for breaking (short breaks of 2–5 minutes) the sedentary time of LTCF residents, twice or 3 times a day. LTCF staff should systematically use simple strategies to stimulate residents to move, such as walking to the lunch/dining hall rather than using wheelchairs for people who are able to ambulate. The risk/benefit of using drugs that reduce patients' activity levels, especially psychotropic drugs, must be systematically evaluated. Physical restraints should be avoided. The use of medical equipment and LTCF architecture should be oriented to optimize residents' mobility. LTCF staff should organize group activities that are motivating and pleasant, such as groups to look after the garden, dance, or walk in green spaces. The use of innovative solutions, such as using animal interventions and new technologies (eg, robots), should be encouraged to increase residents' motivation and pleasure and, then, overall activity levels. 	<ol style="list-style-type: none"> Every resident who has no contraindications must have a personalized exercise program as part of his or her health care plan. Exercise type. The best exercise type is a multicomponent training composed of muscle strength and cardiorespiratory endurance exercises as the core components. Other exercise types, particularly flexibility and balance, should be added to the exercise program whenever possible. Exercise intensity. Moderate-intensity exercises are feasible, effective, and safe. Moderate exercises can be achieved by performing the following: (1) Strength, 1 or 2 sets of exercises, performed at 13–15 repetitions maximum; (2) Aerobic, exercises that noticeably increase heart and respiratory frequency, without generating breathlessness or undue fatigue (scoring 5 or 6 in a 10-point scale of perceived effort). High-intensity exercises can be executed, but may require closer monitoring. Frequency. Twice a week, with an interval of at least 48 hours between sessions. Higher weekly frequency is safe and may be feasible for fitter residents. Duration: 35–45 minutes per session. Lesser durations may be needed during the first weeks of exercise. Longer sessions are feasible for most people.

*Capable of ambulating/rising from a chair with or without human assistance.

Proposed Recommendations to Increase Overall Activity Levels

Although there are no evidence-based guidelines for reducing sedentary time in LTCF settings, task force members recognize the crucial importance of enhancing the overall levels of activity in the daily life of residents. LTCF staff leadership should consider the following:

- (1) To adopt strategies for breaking the sedentary time of LTCF residents. Establishing short breaks (2–5 minutes) 2 or 3 times a day is probably feasible in the LTCF setting.
- (2) To systematically use simple strategies to stimulate residents to move: walking to the lunch/dining hall rather than using wheelchairs for people who are able to ambulate, and organizing events that require residents going out from their rooms.
- (3) To avoid chemical and physical restraints as much as possible because they result in bed and chair rest.
- (4) To optimize the utilization of the LTCF architecture and equipment so as to promote mobility.
- (5) Physical or occupational therapists, or another member of the staff, should organize group activities that are motivating and pleasant. Establishing groups to look after the garden, promoting dancing,⁷⁵ or organizing walks in green spaces⁷⁶ around the LTCF, are examples of potential group activities that can be implemented. Group activities must take into account social affinities among residents, but also residents' interests and preferences to define the most suitable activities to implement.
- (6) To use innovative solutions, such as using animal interventions and new technologies, so as to increase residents' motivation and pleasure and, then, overall activity levels. Animal interventions have been shown to be effective in increasing physical activity in institutionalized older adults.⁷⁷ The use of robots for the institutionalized elderly has shown to decrease feelings of loneliness⁷⁸ and improve participation in activities.⁷⁹ Robots and other technologies (eg, emotional virtual interfaces) have the potential to increase residents' motivation for undertaking (light-intensity) physically demanding activities.

Evidence-Based Exercise Guidelines for a Subgroup of Dependent LTCF Residents: Second Level of Recommendations

Health professionals previously tried to propose some guidance on exercise for LTCF residents.³⁷ However, there is no clear, well-established evidence-based guidelines on exercise for institutionalized older adults currently.

To identify the relevant information for debating on the best exercise regimen in LTCFs, we decided to focus our critical analysis on ADL disability. Indeed, ADL disability is a major issue for the care of institutionalized older people due to its high prevalence in LTCFs,^{1,15,80} its economic burden,^{3,4} and its negative repercussion on residents' quality of life.⁸¹ Moreover, the most robust evidence on the health benefits of exercise for older adults in LTCFs regards its positive impact on ADL function.¹⁰

Procedures

We selected relevant exercise RCTs by proceeding as follows: from the most recent review and meta-analysis of RCTs and cluster-RCTs of exercise for people in LTCF by Crocker and colleagues,¹⁰ we identified the RCTs that met 2 eligibility criteria: (1) had assessed the ability to perform ADLs and (2) used an active exercise intervention; studies of whole-body vibration and staff educational

interventions, if not coupled with an exercise training, were not examined. Then, we extracted data from the original RCTs on exercise type, frequency, intensity, session duration, intervention length, and compliance regarding exercise frequency, as well as on the effects of the intervention on ADL performance. To obtain a value that represents the real frequency of exercise practiced by participants, we multiplied compliance rates by exercise frequency and divided it by 100; data on the real frequency are considered when elaborating the exercise recommendations. We additionally examined the articles classified by Crocker et al¹⁰ as "awaiting assessment" and that were not included in their review. Because Crocker et al¹⁰ performed their last electronic search in August 2011, we performed an updated search from August 2011 to present (searches were performed on October 13, 2015) by using simple key terms (see supplementary materials) in PubMed, the Cochrane Central Register of Controlled Trials, SportDiscus, and PsychInfo databases. This electronic search was intended to incorporate recent RCTs that have investigated the effects of exercise on the ability to perform ADLs in the LTCF setting; a comprehensive systematic review intending to update the study of Crocker et al¹⁰ was out of the scope of this task force. Although examining all RCTs that met the eligibility criteria, we focused our critical analysis on RCTs that found positive effects of exercise on ADL performance, particularly those with more than 100 participants. The rationale for this procedure is related to the fact that Crocker et al¹¹ found that larger RCTs had more conservative results (maybe more realistic results) on the benefits of exercise on the ability to perform ADLs.

After those search procedures, we obtained 32 RCTs that investigated the effects of exercise on ADL performance in LTCF residents; task force members added another exercise RCT meeting the eligibility criteria (Table 2). Almost all 33 RCTs used measures of functional ability as eligibility criteria for the selection of the study population. Ability to ambulate a few meters (with or without human assistance, according to the study) or ability to rise from a chair (without human assistance, with the assistance of 1 or even 2 caregivers, according to the study) were frequent eligibility criteria as well as dependency in basic ADLs (1, 2, or even more ADLs); therefore, the older LTCF residents participating in those RCTs were dependent in 1 or several basic ADLs, but were able to ambulate a few meters and/or to rise from a chair. The baseline levels of functional ability of participants in the exercise group in those 33 RCTs can be characterized by an average Barthel Index of 68 of 100 ($n = 6$ studies,^{24,83,86,102,105,108} varying from 34.0 to 89.6) or 11.9 of 20.0 ($n = 7$ studies,^{40,84,99–101,104,107} varying from 10.1 to 16.1), a functional independence measure (FIM) of 78.1 of 126.0 ($n = 5$ studies,^{24,82,94,96,98} varying from 48.0 to 114.7), or a Rivermead mobility index (RMI) of 5.7 of 15 ($n = 4$ studies,^{84,99–101} varying from 4.9 to 6.1); for all those measurements, higher scores are better. Regarding cognition, participants of those trials had varied levels of cognitive function, transitioning from intact cognition to severe dementia; at least 4 studies^{83,85,86,95} excluded people with dementia and at least 4 other studies^{28,102,106,108} included only people with dementia. Most studies had participants with some degree of cognitive decline. The baseline mini-mental state examination (MMSE) score for the exercise group was, on average, lower than 24 in all^{24,25,27,28,40,82,84,91,94,97,101,102,106–108} but 1 study⁹² that used this assessment tool; the mean MMSE varied from 6.1⁸⁴ to 25⁹² across studies.

Table 2 shows the characteristics of the 33 studies and their exercise interventions. The most represented countries were the United States ($n = 7$); the United Kingdom ($n = 4$); and France, Spain, and the Netherlands ($n = 3$ each). Mean values for the number of study participants was 135.8, mean age on average was 82.5 years, with 74.7% being women. The average exercise program (information for the 2 exercise groups in Deschamps et al⁹¹ and Faber et al⁹² and for the 3 exercise groups in Chin A Paw et al²³ were combined) lasted

Table 2
Study Characteristics and Exercise Interventions

Study	Country	Study Population			ADL Measure	Exercise Intervention					Control Group	
		n	Age*	Sex, % female [†]		Length, wk [‡]	Frequency, weekly [§]	Compliance [§]	Session Duration, min [§]	Intensity		Type
Baum 2003 ⁸²	USA	21	88	75	PPT	25.8	3	80	60	Moderate	Multicomponent (strength + flexibility)	Social activities
Benavent-Caballer 2014 ⁸³	Spain	89	83.9	65.1	Barthel Index	16	3	100	32.5	Light	Strength training	Usual care
Brill 1998 ²¹	USA	16	82	87	ADL score (eg, dressing, transferring)	8	3	93	30	Light-to-moderate	Strength training	Light exercise
Brittle 2009 ⁸⁴	UK	56	84.5	71	RMI	5	2	42.5	50	Light-to-moderate	Multicomponent (balance/coordination + strength + flexibility + aerobic)	Usual care
Cadore 2014 ⁸⁵	Spain	32	91.7		Barthel Index	12	2	90	40	Moderate	Multicomponent (high-speed strength training [for muscle power] + balance/coordination + flexibility)	Light exercise
Chen 2015 ⁸⁶	Taiwan	127	79.1	49.1	Barthel Index	25.8	3	94.5	40	Unclear	Multicomponent (strength + flexibility + aerobics)	Usual care
Chin A Paw 2004 ²³ /Chin A Paw 2006 ⁸⁷	Netherlands	224	81.1	80.2	ADL score (17 activities)	24	2	73 [¶]	52.5	Moderate-to-high	Multicomponent (3 exercise groups. Strength: strength + stretching; Functional: functional activities + stretching; Combined: a combination of the other 2 exercise groups)	Social activities
Conradsson 2010 ⁴⁰ /Rosendahl 2006 ⁸⁸ /Littbrand 2006 ⁸⁹ /Littbrand 2009 ⁹⁰ /Deschamps 2010 ⁹¹	Sweden France	191 160	84.7 82.3	73 71.7	Barthel Index Katz ADL index	13 26	2.2 4 TC 2 CA	72.2 38.8 TC 48.9 CA	45 30 TC 35 CA	Moderate-to-high Light-to-moderate	Multicomponent (balance/coordination + strength + functional training) Multicomponent (TC: Tai chi + muscle reinforcement; CA: balance + strength + flexibility)	Social activities
Dorner 2007 ²⁴	Austria	42	86.8	77	Barthel Index and FIM	10	3	91.8	50	Unclear	Multicomponent (balance/coordination + strength)	Active control
Faber 2006 ⁹²	Netherlands	278	84.9	79	GARS	20	2	88 FW 84 IB	60	Moderate	Multicomponent (IB: adapted Tai Chi + balance; FW: functional training + balance)	Usual care
Gallon 2011 ⁹³	Brazil	19	67	100	Barthel Index	8	3			Light	Flexibility	Social activities
Grönstedt 2013 ⁹⁴	Sweden Denmark Norway	322	85	73.5	FIM	12	4		29.25	Unclear	Multicomponent (balance/coordination + strength + aerobic + functional training)	Usual care
Hsu 2011 ⁹⁵	Canada	44	80	71	NHPPT	4**	3	92 ^{††}	20 ^{††}	Moderate	Multicomponent (balance/coordination + strength) + Wii (bowling)	Multicomponent exercise
Kerse 2008 ²⁹	New Zealand	682	84.3	74	LLFDI	25.8	Daily			Unclear	Functional training (physical activities based on repetitions of ADLs, such as rising from a chair, walking, or repeated transfers)	Social activities
Lazowski 1999 ⁹⁶	Canada	96	80	84	FIM	17.2	3	86	45	Moderate	Multicomponent (balance/coordination + strength + aerobic + flexibility)	Flexibility
Lorenz 2012 ⁹⁷	USA	193	81	64.7	NHPPT	7	5	81	28.4	High	Multicomponent (strength + aerobic)	Usual care
Makita 2006 ⁹⁸	Japan	149	85	100	FIM	12.9	3			Unclear	Multicomponent (balance/coordination + strength + aerobic)	Unclear
McMurdo 1993 ²²	Scotland	49	81	80	Barthel Index	30.1	2	91	45	Light	Multicomponent (strength + flexibility)	Social activities
Meuleman 2000 ²⁵	USA	78	75	12	PADL	6	5		30	Moderate	Multicomponent (strength + aerobic)	Usual care
Morris 1999 ²⁶	USA	468	84.7	79	MDS: Late and early loss ADL and ADL summary	40	5		20	Moderate	Multicomponent (strength + aerobic)	Usual care
Mulrow 1994 ²⁷	USA	194	80	71	Physical Disability Index and Katz ADL	17.2	3	89	30	Moderate	Multicomponent (balance/coordination + strength + flexibility + aerobic + functional training)	Social activities
Rolland 2007 ²⁸	France	134	83	75	Katz ADL	52	2	33	60	Moderate	Multicomponent (balance/coordination + strength + flexibility + aerobic)	Usual care
Sackley 2006 ⁹⁹	UK	118	87	82	RMI and Barthel Index	12	0.28 ^{§§}		30	Unclear	Multicomponent (flexibility + aerobic + functional training)	Usual care
Sackley 2008 ^{100,††}	UK	34	86	88	RMI	4	2		60	Unclear	Multicomponent (balance/coordination + strength + flexibility + aerobic + functional training)	Usual care
Sackley 2009 ¹⁰¹	UK	249	85	74	RMI and Barthel Index	12.9	1.25 ^{§§}		21.5	Moderate	Multicomponent (balance/coordination + strength + flexibility + aerobic + functional training)	Usual care
Santana-Sosa 2008 ¹⁰²	Spain	16	74.5	38	Katz ADL and Barthel Index	12	3	98.9	75	Light	Multicomponent (balance/coordination + strength + flexibility)	Usual care

Seynnes 2004 ¹⁰³	France	27	81.5	Disability index	10	3	99	8	High and light (two exercise groups)	Strength training	Placebo
<i>Tak 2012¹⁰⁴</i>	Netherlands	192	84.6	100	PPT	22	1	30	Light	Multicomponent (aerobic + functional training)	Usual care
<i>Takeuchi 2011¹⁰⁵</i>	Japan	28	85.7	85.7	Barthel Index	4	2	100	Light	Multicomponent (balance/coordination + strength + aerobic)	Multicomponent without aerobics (walking)
<i>Tappen 1994¹⁰⁶</i>	USA	72	84	75	Physical Self-Maintenance Scale and PADL Barthel Index	20	5	150	Unclear	Functional training	Usual care
<i>Tsaih 2012¹⁰⁷</i>	Taiwan	59	76.9	60	Barthel Index	4	3	100	Light	Multicomponent (balance/coordination + aerobic + functional training)	Usual care
<i>Venturelli 2011¹⁰⁸</i>	Italy	24	84	100	Barthel Index and PPT	24	4	93.4	Moderate	Walking	Usual care

CA, cognition-action group; FIM, functional independence measure; FW, functional walking group; GARS, Groningen Activity Restriction Scale; IB, in balance group; LLFDI, Late Life Function and Disability Instrument; MDS, minimum data set; NHPPT, Nursing Home Physical Performance Test; PADL, Physical Activities of Daily Living; PPT, physical performance test; TC, Tai Chi.

Studies showing a positive effect of exercise, compared with controls, on ADL performance (global scores or scores of validated subscales) are italicized.

^aWhen participants' age was provided separately by group, we added all ages and divided by the number of study groups to obtain a mean age.

[†]If % of female provided separately by group, we added all % of female and divided by the number of study groups to obtain a mean. Sex may also have been calculated from the absolute number of males and females in the studies.

[‡]If intervention length was provided in months, we then multiplied the months by 4.3 to obtain "weeks."

[§]If this information was provided as a range, we calculated a mean by adding upper and lower limits and dividing by 2.

^{||}In those studies, intensity was not clearly described by the authors. When possible, we defined the intensity from "cues" provided by the authors. For example, in the study by Baum 2003⁸² rated as *moderate*: "Each exercise was begun with one set of 5 repetitions and gradually progressed to two sets of 10 **as tolerated**. Participants were evaluated weekly for the quality of their movement and number of repetitions with good technique to determine progression of soft weights, balls, or resistance of the therabands."

[¶]Information on the 3 exercise groups (strength training, functional training, and a combination of strength and functional training) were combined because none of these groups improved ADL function compared with controls. This allowed us to obtain a mean value for the 3 groups.

^{**}Before the crossing over of the interventions.

^{††}Information regards only the Wii part of the intervention.

^{‡‡}Although the "mean mobility scores (as measured by the RMI) declined in the control group (n = 16; baseline: 5.9, 6 weeks: 4.75) when compared with the intervention group (n = 17; baseline: 6.1, 6 weeks: 6.2)," the authors did not present any formal statistical analysis.

^{§§}The exact weekly frequency was not provided, but the authors reported the total exposure (number of visits and total duration). From this, we calculated average weekly frequency and session duration.

approximately 16.4 weeks, with a weekly frequency of 2.9 sessions (the “daily” frequency of Kerse et al²⁹ was replaced by 5), a compliance rate of 83.2% (n = 22 RCTs), and a session duration of 42 minutes; real frequency performed (frequency × compliance/100) was 2.3 times per week. The main type of exercise was by far a moderate-intensity multicomponent training (a combination of 2 or more exercise types, such as balance, strength, aerobic, flexibility); the exercise types used across studies were strength (n = 25), balance/coordination (n = 17), aerobic/cardiorespiratory endurance (n = 16), and flexibility (n = 15). None of the trials reported major adverse events (death, important cardiovascular events [eg, heart attack], severely injurious falls, or fractures) related to the exercise intervention; at least 2 studies^{27,28} reported that exercisers fell more than controls, but this difference did not reach statistical significance in any of the trials. It must be, however, highlighted that not all studies clearly reported adverse events.

Fifteen RCTs found a positive effect of exercise on measures of ADL performance (studies italicized in Table 2). The characteristics of these RCTs were similar to those described for the whole sample: the average number of study participants was 124.5, aged 82.8 years, 73.4% women. The average exercise program lasted almost 24 weeks, with a weekly frequency of 3.1 sessions, a compliance rate of 81.7% (n = 12 RCTs), and a session duration of approximately 48 minutes; real frequency performed was 2.5 times per week. The main type of exercise was by far a moderate-intensity multicomponent training; the exercise types more frequently used across studies were strength (n = 12), aerobic/cardiorespiratory endurance (n = 8), flexibility (n = 9), and balance/coordination (n = 6). Focusing on the 7 RCTs in which the exercise intervention was found to improve ADL function and that had more than 100 participants, we obtained the following results: the mean number of study participants was 209, aged approximately 82 years, 72.9% women. The average exercise program lasted almost 27 weeks, with a weekly frequency of 3.1, a compliance rate of 68% (n = 5 RCTs), and a session duration of 34 minutes; real frequency performed was 2.3 times per week. The main type of exercise was a moderate-intensity multicomponent training (all 7 studies used a multicomponent training); the exercise types more used across studies were strength (n = 6), aerobic/cardiorespiratory endurance (n = 6), flexibility (n = 4), and balance/coordination (n = 3).

Proposed Guidelines for Exercise Training

The evidence shows that a moderate-intensity multicomponent training program, composed of strength and aerobic exercises, is effective for improving ADL performance in older adults living in LTCFs; additional benefits may be achieved by integrating balance/coordination and flexibility exercises to the multicomponent training. The proposed guidelines should be applied for institutionalized older adults who are dependent in basic ADLs, but still able to ambulate and/or to rise from a chair (with or without human assistance), including people who had cognitive decline and even dementia; indeed, the evidence used to build these guidelines was extracted from populations with such characteristics, as described previously. Moreover, these recommendations should be applied for overall stable patients. Patients in the end of life, bed-bound, or with any unstable conditions precluding exercise should be examined on a case-by-case basis and should have a more personalized intervention, if appropriate. To be effective on a long-term basis, the exercise program must be continuously adapted to individuals' capacity and must be organized as a progressive challenge. Residents should be stimulated to comply with these guidelines and clear strategies to “reengage” those who quit exercise classes should be defined by the LTCF staff leadership.

The task force group makes the following recommendations as the minimum exercise training to be provided for this subgroup of LTCF residents.

Exercise Type

The type of exercise should be multicomponent training. Exercises designed for improving muscle strength and cardiorespiratory endurance must be the essential part of the multicomponent training. Strength training may be performed through weight-bearing exercises and using free weights (the use of machines is less usual in the LTCF setting, even though it is feasible). Special attention must be paid to strengthening the lower body. The aerobic aspect may traditionally be implemented through walking (in continuum or by walk bouts); using circuit training, with walk bouts interposed between other types of exercises, may also contribute to achieve aerobic endurance. Other exercise types, particularly flexibility and balance, should be added to the exercise program whenever possible. For example, active range-of-motion exercises may be integrated in the warm-up phase, and stretching may be part of the cool-down phase of the training. To increase strength and, then, self-confidence and mastery, several sessions of strength training may be needed before introducing balance/coordination activities, particularly for very vulnerable people. Specific exercises may be integrated so as to train people for the execution of actions that are strongly associated with falls, particularly standing quietly, sitting down or lowering, and initiation of walking¹⁰⁹; adding a component of muscle power (prioritizing the speed of muscle contraction) may be integrated in such an exercise training. Walking forward with direction changes may also contribute to reduce falls, as walking forward is the most common activity at the moment of fall.¹⁰⁹

Intensity

Exercise intensity should be moderate. Moderate intensity can be achieved in the following ways for the different types of exercise.

Strength

Strength training included 1 or 2 sets of exercises, performed at 13 to 15 repetitions maximum. The number of exercises will be limited by time availability and residents' exercise capacity; the ideal number of exercises would be 8 to 10. Low-intensity exercises performed at up to 20 repetitions maximum may be required, particularly in the first weeks of the exercise program. High-intensity exercise (ie, exercises performed at 8–12 repetitions maximum) can be executed, but it may require closer monitoring.⁸⁹

Aerobic endurance

Aerobic endurance is achieved through exercises that noticeably increase heart rate and respiration, without generating breathlessness or undue fatigue. It can be easily monitored through self-report as recommended by the American College of Sports Medicine/American Heart Association: “On a 10-point scale, where sitting is 0 and all-out effort is 10, moderate-intensity activity is a 5 or 6 and produces noticeable increases in heart rate and breathing.”¹³ Attention must be paid for people with important cognitive impairment, who are unable to identify moderate intensity and to communicate potential acute signs, such as pain. Light-intensity exercises appear to provide some benefits and should be envisaged for very vulnerable people and at the start of any exercise program (during the first weeks of exercise). High-intensity exercise can be performed, but it may require a closer monitoring; on the same self-reported scale as above, “vigorous-intensity activity is a 7 or 8 and produces large increases in heart rate and breathing.”¹³

Flexibility

Active range-of-motion and stretch exercises should be performed during 10 to 30 seconds per exercise; the number of exercises will be limited by time constraints. Although assessing the intensity of flexibility exercises is very difficult, asking residents to perform range-of-motion and stretching exercises to the maximum of their capacity without reaching painful levels is probably feasible and safe. Attention must be paid to people who are unable to communicate painful feelings.

Balance

Establishing the intensity of balance exercises is a challenge because there are no validated instruments developed to this purpose.¹¹⁰ A progressive increase in the difficulty for executing static (eg, semi-tandem, tandem, single-leg stand), but mainly dynamic balance (eg, walking in the line, tandem walking in the line; walking with changing directions slowly and, then, faster) is required. Exercises that reduce sensory input (eg, standing with eyes closed)¹² also can be used.

Frequency

Exercises should be performed twice a week. More than twice a week can be a difficult target for most LTCF residents, and doing less than that can be ineffective.⁸⁷ An interval of at least 48 hours between sessions should be respected. An exercise program with 3 or more weekly sessions is, however, safe and may be feasible for fitter residents.

Duration

Duration of exercise should be between 35 and 45 minutes per session. This session duration range gives enough time for the essential components of the exercise training (ie, strength and aerobic training), and still leaves some time for other exercises, such as flexibility and balance. Lesser durations may be needed for very vulnerable people and during the first weeks of exercise. Longer sessions (preferentially, not longer than 60 minutes) are, however, feasible for most people.

A Typical Exercise Session for Older People in LTCFs

The following is an example of how a 45-minute session of exercise could be divided:

- 4 minutes of warm-up. Range-of-motion exercises (for example, for the wrists, shoulders, hip, knees, and ankles), followed by light walking.
- 8 minutes balance/coordination. Standing balance with increasing difficulty (eg, narrowing the base of support); activities of bodyweight shift; walking forward with changing directions; walking along a straight line (forward, backward, and sideways).
- 15 minutes strength. Thirteen to 15 repetitions maximum of chair rises, with increasing difficulty (eg, emphasizing speed of movement); different theraband exercises for the upper body and trunk; calf weights for knee extension and flexion or weighted belts for functional lower-limb strength exercises.
- 15 minutes aerobic. Five 3-minute bouts of walking interposed between 2 strength exercises and/or between 2 balance/coordination exercises.
- 3 minutes cool down. Very light walking followed by a few stretching exercises.

Implementing the Recommendations

To promote the implementation of the proposed recommendations in the real life of LTCFs, it is crucial to take into account residents' desires, preferences, beliefs, and attitudes toward physical activities and exercises. For instance, taking into account a subject's self-efficacy (ie, one's belief that he or she is capable of performing the goal-directed behavior) to define challenging but feasible goals (with the resident whenever possible) are essential factors to consider when designing strategies to increase/maintain optimal levels of physical activity and exercise. Other important aspects for the successful implementation of the recommendations are related to promoting social support (eg, by doing physical activities and exercise in group), and providing a stimulating environment (eg, by using different equipment with different colors: free weights, balls, carpets with different consistencies, by using music during the exercise sessions, and even implementing dual-task training) for LTCF residents. Building awareness of residents' family members and primary health care providers about the importance of physical activity and exercise for older adults as well as about their role in keeping the resident physically active is a key aspect for the implementation of our recommendations.

Moreover, LTCF organizational aspects are as important as the recommendations themselves. Even though it is out of the scope of this task force to deeply debate about LTCF organizational aspects, task force members recognize that strategies that may favor a successful exercise implementation in LTCFs should be discussed at the facility and health care system levels. At the facility level, LTCF leadership staff may discuss and implement different strategies that include (but are not limited to) the following: involving residents' family and primary health care provider in the "exercise strategy" to increase residents' adherence to exercise; adding exercise programs in the personalized health care plan of all residents who do not have any contraindication for exercising (every resident who has no contraindications must have a personalized exercise program); defining strategies for providing incentives to residents who exercise regularly, for example, by rewarding residents who increased or maintained high attendance rates in exercise sessions; defining an "exercise referent" person among LTCF staff, as it already exists for pain care; and providing sufficient training to LTCF staff and volunteers in how to conduct exercise sessions. To optimize the potential health benefits of physical activity and exercise in LTCF residents, having extra people in the LTCF is required. The training level in terms of knowledge and experience in physical activity of the extra people can vary from highly trained professionals to sufficiently trained volunteers. Although the ideal picture is that exercise training sessions in the LTCF should be led by professionals from the field of exercise sciences, and all efforts must be undertaken toward this ideal picture, an important number of facilities (maybe most of them) will not have the required resources for that; in those LTCFs, health care providers, other than exercise science professionals, and/or volunteers will be in charge of conducting exercise sessions, needing thus to achieve minimum levels of knowledge and expertise on exercise.

In a transition level, which would ask for efforts from both facility and health care systems, strategies that build awareness about the potential role of the LTCF as a site for health care prevention through exercise should be promoted; for example, by taking advantage of the already existing space and staff expertise in LTCFs, facilities may open their doors and invite the whole community to exercise in the LTCF, with LTCF residents. Other innovative solutions, such as integrating volunteers, including high school or college students, into the strategies of the LTCF so as to increase both residents' daily life physical activity (eg, walking or gardening with volunteers) and their adherence to exercise sessions (eg, exercising with volunteers) should also be welcome; volunteer-led exercise interventions have already been

tested and provided positive results on exercise adherence (participants completed on average 94% of the proposed volunteer-led sessions) and on clinical outcomes (eg, ADL performance).⁸⁶

At the level of health care systems, policies that facilitate the implementation of exercise in the LTCF setting should be debated and implemented. For example, establishing the successful organization of exercise sessions in the LTCF (for example, by conducting exercises according to the guidelines proposed herein) as a quality indicator of care and/or creating a distinguished label (“LTCF fit-friendly”) for LTCFs that organize exercise successfully, and defining ways to pay/reimburse the expenses related to the exercise program, would certainly help in promoting exercise in LTCFs.

Conclusions and Perspectives

Scientific evidence about the effects of physical activity and exercise on the health of older adults in LTCFs has increased in the past decade. Establishing strategies to increase activity levels in daily life as well as minimal standards in terms of exercise regimen is important for the guidance of professionals working in this setting.

Task force members purposefully covered a very large spectrum of recommendations from overall strategies to increase physical activity levels in the daily life of LTCF residents to more precise guidelines on exercise training to a well-defined subgroup of LTCF residents. Our recommendations should not be used as a one-size-fits-all approach. Instead, they should be used as a flexible framework, as the most appropriate strategies to increase residents' daily life physical activity will vary across facilities, and the different parts of the exercise regimen (type, frequency, intensity, duration) should be adapted according to residents' capacities, needs, and desires. It is important to highlight that, currently, LTCF residents appear to engage in less exercise than the recommendations proposed herein, as showed in an observational cross-sectional study in which only 10% of nursing home residents exercised at least twice a week.¹¹¹

To confirm that our recommendations in terms of overall physical activity and exercise training are appropriate and effective for institutionalized older adults, further studies are needed. Well-designed and large RCTs are particularly welcome to examine the effectiveness of the exercise recommendations on different relevant outcomes for LTCF residents, such as BPSD, falls, malnutrition, and quality of life, as well as on staff-related outcomes, such as reductions of staff burnout. Cost-effectiveness analysis is also required to strengthen our recommendations.

Supplementary Data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.jamda.2016.01.021>.

References

- De Souto Barreto P, Lapeyre-Mestre M, Mathieu C, et al. A multicentric individually-tailored controlled trial of education and professional support to nursing home staff: Research protocol and baseline data of the IQUARE study. *J Nutr Health Aging* 2013;17:173–178.
- United Nations, Department of Economic and Social Affairs, Population Division. *World Population Ageing 2013*. New York (NY): United Nations publication; 2013;ST/ESA/SER.A/348.
- Pickard L, Comas-Herrera A, Costa-Font J, et al. Modelling an entitlement to long-term care services for older people in Europe: Projections for long-term care expenditure to 2050. *J Eur Soc Policy* 2007;17:33–48.
- De Meijer CAM, Majer IM, Koopmanschap MA, Van Baal PH. Forecasting lifetime and aggregate long-term care spending: Accounting for changing disability patterns. *Med Care* 2012;50:722–729.
- Lubitz J, Cai L, Kramarow E, Lentzner H. Health, life expectancy, and health care spending among the elderly. *N Engl J Med* 2003;349:1048–1055.
- Wübker A, Zwakhalen SM, Challis D, et al. Costs of care for people with dementia just before and after nursing home placement: Primary data from eight European countries. *Eur J Health Econ* 2015;16:689–707.
- World Health Organization. *World report on ageing and health*. Geneva: WHO Press; 2015. p. 247.
- Tak E, Kuiper R, Chorus A, Hopman-Rock M. Prevention of onset and progression of basic ADL disability by physical activity in community dwelling older adults: A meta-analysis. *Ageing Res Rev* 2013;12:329–338.
- Tolson D, Rolland Y, Andrieu S, et al. International Association of Gerontology and Geriatrics: A global agenda for clinical research and quality of care in nursing homes. *J Am Med Dir Assoc* 2011;12:184–189.
- Crocker T, Forster A, Young J, et al. Physical rehabilitation for older people in long-term care. *Cochrane Database Syst Rev* 2013;(2):CD004294.
- Crocker T, Young J, Forster A, et al. The effect of physical rehabilitation on activities of daily living in older residents of long-term care facilities: Systematic review with meta-analysis. *Age Ageing* 2013;42:682–688.
- American College of Sports Medicine, Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, et al. American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc* 2009;41:1510–1530.
- Nelson ME, Rejeski WJ, Blair SN, et al. Physical activity and public health in older adults: Recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39:1435–1445.
- World Health Organization. *Global recommendations on physical activity for health*. Geneva: WHO Press; 2010. p. 58.
- Rolland Y, Abellan van Kan G, Hermabessiere S, et al. Descriptive study of nursing home residents from the REHPA network. *J Nutr Health Aging* 2009;13:679–683.
- Morley JE, Caplan G, Cesari M, et al. International survey of nursing home research priorities. *J Am Med Dir Assoc* 2014;15:309–312.
- Forbes D, Forbes SC, Blake CM, et al. Exercise programs for people with dementia. *Cochrane Database Syst Rev* 2015;(4):CD006489.
- Pitkälä K, Savikko N, Poysti M, et al. Efficacy of physical exercise intervention on mobility and physical functioning in older people with dementia: A systematic review. *Exp Gerontol* 2013;48:85–93.
- de Souto Barreto P, Demougeot L, Pillard F, et al. Exercise training for managing behavioral and psychological symptoms in people with dementia: A systematic review and meta-analysis. *Ageing Res Rev* 2015;24:274–285.
- Buettner LL, Ferrario J. Therapeutic recreation-nursing team: A therapeutic intervention for nursing home residents with dementia. *Annual in Therapeutic Recreation* 1997–1998;7:21–28.
- Brill P, Jensen R, Koltyn K, et al. The feasibility of conducting a group-based progressive strength training program in residents of a multi-level care facility. *Act Adapt Aging* 1998;22:53–63.
- McMurdo MET, Rennie LM. A controlled trial of exercise by residents of old people's homes. *Age Ageing* 1993;22:11–15.
- Chin A Paw MJ, van Poppel MNM, Twisk JWR, van Mechelen W. Effects of resistance and all-round, functional training on quality of life, vitality and depression of older adults living in long-term care facilities: A 'randomized' controlled trial. *BMC Geriatr* 2004;4:5.
- Dorner T, Kranz A, Zettl-Wiedner K, et al. The effect of structured strength and balance training on cognitive function in frail, cognitive impaired elderly long-term care residents. *Ageing Clin Exp Res* 2007;19:400–405.
- Meuleman JR, Brechue WF, Kubilis PS, Lowenthal DT. Exercise training in the debilitated aged: Strength and functional outcomes. *Arch Phys Med Rehabil* 2000;81:312–318.
- Morris JN, Fiatarone M, Kiely DK, et al. Nursing rehabilitation and exercise strategies in the nursing home. *J Gerontol A Biol Sci Med Sci* 1999;54:M494–M500.
- Mulrow CD, Gerety MB, Kanten D, et al. A randomized trial of physical rehabilitation for very frail nursing home residents. *JAMA* 1994;271:519–524.
- Rolland Y, Pillard F, Klapszczyk A, et al. Exercise program for nursing home residents with Alzheimer's disease: A 1-year randomized, controlled trial. *J Am Geriatr Soc* 2007;55:158–165.
- Kerse N, Peri K, Robinson E, et al. Does a functional activity programme improve function, quality of life, and falls for residents in long term care? Cluster randomised controlled trial. *BMJ* 2008;337:a1445.
- Underwood M, Lamb SE, Eldridge S, et al. Exercise for depression in elderly residents of care homes: A cluster-randomised controlled trial. *Lancet* 2013;382:41–49.
- Cameron ID, Gillespie LD, Robertson MC, et al. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst Rev* 2012;(12):CD005465.
- Gillespie LD, Robertson MC, Gillespie WJ, et al. Interventions for preventing falls in older people living in the Community. *Cochrane Database Syst Rev* 2012;(9):CD007146.
- El-Khoury F, Cassou B, Charles MA, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: Systematic review and meta-analysis of randomised controlled trials. *BMJ* 2013;347:f6234.
- Cadore EL, Rodríguez-Mañas L, Sinclair A, Izquierdo M. Effects of different exercise interventions on risk of falls, gait ability, and balance in physically frail older adults: A systematic review. *Rejuvenation Res* 2013;16:105–114.
- Kojima G. Prevalence of frailty in nursing homes: A systematic review and meta-analysis. *J Am Med Dir Assoc* 2015;16:940–945.
- Luo H, Lum TYS, Wong GHY, et al. Predicting adverse health outcomes in nursing homes: A 9-year longitudinal study and development of the FRAIL-

- Minimum Data Set (MDS) Quick Screening Tool [published online ahead of print October 23, 2015]. *J Am Med Dir Assoc* <http://dx.doi.org/10.1016/j.jamda.2015.09.006>.
37. Weening-Dijksterhuis E, de Greef MH, Scherder EJ, et al. Frail institutionalized older persons: A comprehensive review on physical exercise, physical fitness, activities of daily living, and quality-of-life. *Am J Phys Med Rehabil* 2011;90:156–168.
 38. Lee LY, Lee DT, Woo J. Tai chi and health-related quality of life in nursing home residents. *J Nurs Scholarsh* 2009;41:35–43.
 39. Mihalko SL, McAuley E. Strength training effects on subjective well-being and physical function in the elderly. *J Aging Phys Act* 1996;4:56–68.
 40. Conradsson M, Littbrand H, Lindelof N, et al. Effects of a high-intensity functional exercise programme on depressive symptoms and psychological well-being among older people living in residential care facilities: A cluster-randomized controlled trial. *Aging Ment Health* 2010;14:565–576.
 41. Giné-Garriga M, Roqué-Fíguls M, Coll-Planas L, et al. Physical exercise interventions for improving performance-based measures of physical function in community-dwelling, frail older adults: A systematic review and meta-analysis. *Arch Phys Med Rehabil* 2014;95:753–769.e3.
 42. Blake H, Mo P, Malik S, Thomas S. How effective are physical activity interventions for alleviating depressive symptoms in older people? A systematic review. *Clin Rehabil* 2009;23:873–887.
 43. Chin A Paw MJM, van Poppel MNM, van Mechelen W. Effects of resistance and functional-skills training on habitual activity and constipation among older adults living in long-term care facilities: A randomized controlled trial. *BMC Geriatr* 2006;6:9.
 44. Ikezoe T, Asakawa Y, Shima H, et al. Daytime physical activity patterns and physical fitness in institutionalized elderly women: An exploratory study. *Arch Gerontol Geriatr* 2013;57:221–225.
 45. Tanner RE, Bruncker LB, Agergaard J, et al. Age-related differences in lean mass, protein synthesis and skeletal muscle markers of proteolysis after bed rest and exercise rehabilitation. *J Physiol* 2015;593:4259–4273.
 46. Wall BT, Dirks ML, van Loon LJC. Skeletal muscle atrophy during short-term disuse: Implications for age-related sarcopenia. *Ageing Res Rev* 2013;12:898–906.
 47. Senior HE, Henwood TR, Beller EM, et al. Prevalence and risk factors of sarcopenia among adults living in nursing homes. *Maturitas* 2015;82:418–423.
 48. Smoliner C, Sieber CC, Wirth R. Prevalence of sarcopenia in geriatric hospitalized patients. *J Am Med Dir Assoc* 2014;15:267–272.
 49. Yalcin A, Aras S, Atmis V, et al. Sarcopenia prevalence and factors associated with sarcopenia in older people living in a nursing home in Ankara Turkey [published online ahead of print August 5, 2015]. *Geriatr Gerontol Int* <http://dx.doi.org/10.1111/ggi.12570>.
 50. Landi F, Liperoti R, Fusco D, et al. Sarcopenia and mortality among older nursing home residents. *J Am Med Dir Assoc* 2012;13:121–126.
 51. Cesari M, Fielding R, Bénichou O, et al. Pharmacological interventions in frailty and sarcopenia: Report by the International Conference on Frailty and Sarcopenia Research Task Force. *J Frailty Aging* 2015;4:114–120.
 52. Liu CJ, Latham NK. Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev* 2009;(3):CD002759.
 53. Tieland M, Dirks ML, van der Zwaluw N, et al. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: A randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 2012;13:713–719.
 54. Landi F, Russo A, Bernabei R. Physical activity and behavior in the elderly: A pilot study. *Arch Gerontol Geriatr Suppl* 2004;9:235–241.
 55. US Department of Health and Human Services Centers for Disease Control and Prevention. Quick stats: Ten most common chronic conditions among persons living in residential care facilities—national survey of residential care facilities, United States, 2010. *MMWR Morb Mortal Wkly Rep* 2012;61:603.
 56. Moore KL, Boscardin WJ, Steinman MA, Schwartz JB. Patterns of chronic comorbid medical conditions in older residents of US nursing homes: Differences between the sexes and across the agespan. *J Nutr Health Aging* 2014;18:429–436.
 57. Pescatello LS, MacDonald HV, Lamberti L, Johnson BT. Exercise for hypertension: A prescription update integrating existing recommendations with emerging research. *Curr Hypertens Rep* 2015;17:87.
 58. American College of Sports Medicine position stand. Exercise for patients with coronary artery disease. *Med Sci Sports Exerc* 1994;26:i–v.
 59. Gielen S, Laughlin MH, O'Conner C, Duncker DJ. Exercise training in patients with heart disease: Review of beneficial effects and clinical recommendations. *Prog Cardiovasc Dis* 2015;57:347–355.
 60. Focht BC. Effectiveness of exercise interventions in reducing pain symptoms among older adults with knee osteoarthritis: A review. *J Aging Phys Act* 2006;14:212–235.
 61. Giangregorio LM, Papaioannou A, Macintyre NJ, et al. Too fit to fracture: Exercise recommendations for individuals with osteoporosis or osteoporotic vertebral fracture. *Osteoporos Int* 2014;25:821–835.
 62. Giangregorio LM, McGill S, Wark JD, et al. Too fit to fracture: Outcomes of a Delphi consensus process on physical activity and exercise recommendations for adults with osteoporosis with or without vertebral fractures. *Osteoporos Int* 2015;26:891–910.
 63. Colberg SR, Albright AL, Blissmer BJ, et al. Exercise and type 2 diabetes: American College of Sports Medicine and the American Diabetes Association: Joint position statement. *Exercise and type 2 diabetes. Med Sci Sports Exerc* 2010;42:2282–2303.
 64. Billinger SA, Arena R, Bernhardt J, et al. Physical activity and exercise recommendations for stroke survivors: A statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2014;45:2532–2553.
 65. Mansoubi M, Pearson N, Clemes SA, et al. Energy expenditure during common sitting and standing tasks: Examining the 1.5 MET definition of sedentary behaviour. *BMC Public Health* 2015;15:516.
 66. van der Ploeg HP, Chey T, Korda RJ, et al. Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med* 2012;172:494–500.
 67. Grøntved A, Hu FB. Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: A meta-analysis. *JAMA* 2011;305:2448–2455.
 68. Chau JY, Grunseit AC, Chey T, et al. Daily sitting time and all-cause mortality: A meta-analysis. *PLoS One* 2013;8:e80000.
 69. Keogh JW, Senior H, Beller EM, Henwood T. Prevalence and risk factors for low habitual walking speed in nursing home residents: An observational study. *Arch Phys Med Rehabil* 2015;96:1993–1999.
 70. De Souto Barreto P. Non-pharmacological interventions in the nursing home setting: Does it make any sense to struggle against sedentary behavior among residents? *J Nursing Home Res* 2015;1:52–54.
 71. Sparling PB, Howard BJ, Dunstan DW, Owen N. Recommendations for physical activity in older adults. *BMJ* 2015;350:h100.
 72. Slaughter SE, Estabrooks CA. Optimizing the mobility of residents with dementia: A pilot study promoting healthcare aide uptake of a simple mobility innovation in diverse nursing home settings. *BMC Geriatr* 2013;13:110.
 73. Baert V, Gorus E, Calleeuw K, et al. An administrator's perspective on the organization of physical activity for older adults in long-term care facilities. *J Am Med Dir Assoc* 2016;17:75–84.
 74. Baert V, Gorus E, Guldemont N, et al. Physiotherapists' perceived motivators and barriers for organizing physical activity for older long-term care facility residents. *J Am Med Dir Assoc* 2015;16:371–379.
 75. Vankova H, Holmerova I, Machacova K, et al. The effect of dance on depressive symptoms in nursing home residents. *J Am Med Dir Assoc* 2014;15:582–587.
 76. Moran M, Van Cauwenberg J, Hercky-Linnewiel R, et al. Understanding the relationships between the physical environment and physical activity in older adults: A systematic review of qualitative studies. *Int J Behav Nutr Phys Act* 2014;11:79.
 77. Friedmann E, Galik E, Thomas SA, et al. Evaluation of a pet-assisted living intervention for improving functional status in assisted living residents with mild to moderate cognitive impairment: A pilot study. *Am J Alzheimers Dis Other Demen* 2015;30:276–289.
 78. Robinson H, Macdonald B, Kerse N, Broadbent E. The psychosocial effects of a companion robot: A randomized controlled trial. *J Am Med Dir Assoc* 2013;14:661–667.
 79. Sung H-C, Chang S-M, Chin M-Y, Lee W-L. Robot-assisted therapy for improving social interactions and activity participation among institutionalized older adults: A pilot study. *Asia Pac Psychiatry* 2015;7:1–6.
 80. Rolland Y, Andrieu S, Crochard A, et al. Psychotropic drug consumption at admission and discharge of nursing home residents. *J Am Med Dir Assoc* 2012;13:407.e7–407.e12.
 81. Andersen CK, Wittrup-Jensen KU, Lolk A, et al. Ability to perform activities of daily living is the main factor affecting quality of life in patients with dementia. *Health Qual Life Outcomes* 2004;2:52.
 82. Baum EE, Jarjoura D, Polen AE, et al. Effectiveness of a group exercise program in a long-term care facility: A randomized pilot trial. *J Am Med Dir Assoc* 2003;4:74–80.
 83. Benavent-Caballer V, Rosado-Calatayud P, Segura-Ortí E, et al. Effects of three different low-intensity exercise interventions on physical performance, muscle CSA and activities of daily living: A randomized controlled trial. *Exp Gerontol* 2014;58:159–165.
 84. Brittle N, Patel S, Wright C, et al. An exploratory cluster randomized controlled trial of group exercise on mobility and depression in care home residents. *Clin Rehabil* 2009;23:146–154.
 85. Cadore EL, Casas-Herrero A, Zambom-Ferraresi F, et al. Multicomponent exercises including muscle power training enhance muscle mass, power output, and functional outcomes in institutionalized frail nonagenarians. *Age (Dordr)* 2014;36:773–785.
 86. Chen K-M, Li C-H, Chang Y-H, et al. An elastic band exercise program for older adults using wheelchairs in Taiwan nursing homes: A cluster randomized trial. *Int J Nurs Stud* 2015;52:30–38.
 87. Chin A Paw MJM, van Poppel MNM, Twisk JWR, van Mechelen W. Once a week not enough, twice a week not feasible? A randomised controlled exercise trial in long-term care facilities [ISRCTN87177281]. *Patient Educ Couns* 2006;63:205–214.
 88. Rosendahl E, Lindelöf N, Littbrand H, et al. High-intensity functional exercise program and protein-enriched energy supplement for older persons dependent in activities of daily living: A randomised controlled trial. *Aust J Physiother* 2006;52:105–113.
 89. Littbrand H, Rosendahl E, Lindelöf N, et al. A high-intensity functional weight-bearing exercise program for older people dependent in activities of daily living and living in residential care facilities: Evaluation of the applicability with focus on cognitive function. *Phys Ther* 2006;86:489–498.

90. Littbrand H, Lundin-Olsson L, Gustafson Y, Rosendahl E. The effect of a high-intensity functional exercise program on activities of daily living: A randomized controlled trial in residential care facilities. *J Am Geriatr Soc* 2009; 57:1741–1749.
91. Dechamps A, Diolez P, Thiaudière E, et al. Effects of exercise programs to prevent decline in health-related quality of life in highly deconditioned institutionalized elderly persons: A randomized controlled trial. *Arch Intern Med* 2010;170:162–169.
92. Faber MJ, Bosscher RJ, Chin A Paw MJ, van Wieringen PC. Effects of exercise programs on falls and mobility in frail and pre-frail older adults: A multi-center randomized controlled trial. *Arch Phys Med Rehabil* 2006;87:885–896.
93. Gallon D, Rodacki ALF, Hernandez SG, et al. The effects of stretching on the flexibility, muscle performance and functionality of institutionalized older women. *Braz J Med Biol Res* 2011;44:229–235.
94. Grönstedt H, Frändin K, Bergland A, et al. Effects of individually tailored physical and daily activities in nursing home residents on activities of daily living, physical performance and physical activity level: A randomized controlled trial. *Gerontology* 2013;59:220–229.
95. Hsu JK, Thibodeau R, Wong SJ, et al. A « Wii » bit of fun: The effects of adding Nintendo Wii(®) Bowling to a standard exercise regimen for residents of long-term care with upper extremity dysfunction. *Physiother Theory Pract* 2011;27:185–193.
96. Lazowski DA, Ecclestone NA, Myers AM, et al. A randomized outcome evaluation of group exercise programs in long-term care institutions. *J Gerontol A Biol Sci Med Sci* 1999;54:M621–M628.
97. Lorenz RA, Gooneratne N, Cole CS, et al. Exercise and social activity improve everyday function in long-term care residents. *Am J Geriatr Psychiatry* 2012; 20:468–476.
98. Makita M, Nakadaira H, Yamamoto M. Randomized controlled trial to evaluate effectiveness of exercise therapy (Takizawa Program) for frail elderly. *Environ Health Prev Med* 2006;11:221–227.
99. Sackley C, Wade DT, Mant D, et al. Cluster randomized pilot controlled trial of an occupational therapy intervention for residents with stroke in UK care homes. *Stroke* 2006;37:2336–2341.
100. Sackley CM, Rodriguez NA, van den Berg M, et al. A phase II exploratory cluster randomized controlled trial of a group mobility training and staff education intervention to promote urinary continence in UK care homes. *Clin Rehabil* 2008;22:714–721.
101. Sackley CM, van den Berg ME, Lett K, et al. Effects of a physiotherapy and occupational therapy intervention on mobility and activity in care home residents: A cluster randomised controlled trial. *BMJ* 2009;339:b3123.
102. Santana-Sosa E, Barriopedro MI, López-Mojares LM, et al. Exercise training is beneficial for Alzheimer's patients. *Int J Sports Med* 2008;29:845–850.
103. Seynnes O, Fiatarone Singh MA, Hue O, et al. Physiological and functional responses to low-moderate versus high-intensity progressive resistance training in frail elders. *J Gerontol A Biol Sci Med Sci* 2004;59: 503–509.
104. Tak ECPM, van Hespren A, van Dommelen P, Hopman-Rock M. Does improved functional performance help to reduce urinary incontinence in institutionalized older women? A multicenter randomized clinical trial. *BMC Geriatr* 2012;12:51.
105. Takeuchi R, Hatano Y, Yamasaki M. The influence of different exercise intervention programs on changes in quality of life and Activity of Daily Living levels among geriatric nursing home residents. *J Phys Ther Sci* 2011;23: 133–136.
106. Tappen RM. The effect of skill training on functional abilities of nursing home residents with dementia. *Res Nurs Health* 1994;17:159–165.
107. Tsaih P-L, Shih Y-L, Hu M-H. Low-intensity task-oriented exercise for ambulation-challenged residents in long-term care facilities: A randomized, controlled trial. *Am J Phys Med Rehabil* 2012;91:616–624.
108. Venturelli M, Scarsini R, Schena F. Six-month walking program changes cognitive and ADL performance in patients with Alzheimer. *Am J Alzheimers Dis Other Demen* 2011;26:381–388.
109. Robinovitch SN, Feldman F, Yang Y, et al. Video capture of the circumstances of falls in elderly people residing in long-term care: An observational study. *Lancet* 2013;381:47–54.
110. Farlie MK, Robins L, Keating JL, et al. Intensity of challenge to the balance system is not reported in the prescription of balance exercises in randomised trials: A systematic review. *J Physiother* 2013;59:227–235.
111. De Souto Barreto P, Demougeot L, Vellas B, Rolland Y. How much exercise are older adults living in nursing homes doing in daily life? A cross-sectional study. *J Sports Sci* 2015;33:116–124.